

IFT-P.42/97

hep-ph/yymmddd

July 1997

The electromagnetic gauge invariance in $SU(3)_L \otimes U(1)_N$ models of electroweak unification reexamined

V. Pleitez^(a) and M. D. Tonasse^(b)

^(a) *Instituto de Física Teórica*

Universidade Estadual Paulista

Rua Pamplona 145

01405-900 – São Paulo, SP

Brazil

^(b) *Instituto de Física da Universidade do Estado do Rio de Janeiro*

Rua São Francisco Xavier 524

20550-013 – Rio de Janeiro, RJ

Brazil

(February 1, 2008)

Abstract

Two models with $SU(3)_C \otimes SU(3)_L \otimes U(1)_N$ gauge symmetry are considered. We show that the masslessness of the photon does not prevent the neutrinos from acquiring Majorana masses. That is, there is no relation between the VEVs of Higgs fields and the electromagnetic gauge invariance contrary to what has been claimed recently.

PACS numbers: 12.60.-i; 12.60.Cn; 12.60.Fr

In model building of the electroweak interactions among elementary particles it is usually assumed that any neutral scalar field can get a Vacuum Expectation Value (VEV). This assures the conservation of the electric charge and the Abelian gauge invariance needed for maintaining the photon massless.

In general the pattern of the symmetry breaking is not arbitrary but depends on the structure of the theory, in particular on the group representation content of the scalar fields [1]. For instance consider the case of $SU(2)$ gauge theory with a complex doublet of scalar fields φ . In the unitary gauge the vacuum expectation value (VEV) is $\langle\varphi\rangle = (0, v)^T$. Then, the original $SU(2)$ symmetry is completely broken. On the other hand, if we have instead a real triplet ϕ with $\langle\phi\rangle = (0, 0, v)^T$ (and we can always chose this vacuum state), the gauge symmetry $SU(2)$ will be broken down to a residual $U(1)$ gauge symmetry with one massless vector boson remaining. In general to break the $SU(n)$ symmetry completely we need $n - 1$ sets of vector representations [1]. Charged scalars will spoil the electromagnetic gauge invariance in a trivial way. Hence, all discussions go around neutral scalar bosons which can, in principle, gain a VEV. It is not obvious that with the proliferation of Higgs fields at least one massless gauge boson, which can be identified as a photon, remains. So, the general recipe for model building [2] has to be checked case by case. For instance, in the case of $SU(2)\otimes U(1)$ theory one doublet with vacuum state $(0, v)^T$ chooses the W^3 direction so, if we add a triplet with vacuum state $(0, v, 0)^T$ the photon will gain a mass. However, this is only an apparent effect since it is always possible to choose the vacuum state of the triplet $(0, 0, v)^T$ in such a way that both vacuum states, the doublet's and the triplet's ones, are always in the same direction. Although this is always possible with a minimal set of scalar multiplets, there are models in which the scalars needed for breaking down the symmetry are not enough to generate all fermion masses. New scalar multiplets are then necessary to generate the fermionic masses and to have still the appropriate symmetry breaking. Hence, we can wonder if in this case that feature will continue to be valid. In fact, it has been claimed by Özer [3] and Pisano *et al.* [4], that in some models based on $SU(3)_C \otimes SU(3)_L \otimes U(1)_N$ symmetry (3-3-1 models) some neutral components of one sextet

of scalar fields cannot gain a VEV since in this case it will break the electromagnetic $U(1)$ gauge symmetry generating a massive photon. The goal of this paper is to show that this is not true in any of those models.

If we consider only leptons with electric charge 0 and ± 1 , the extensions of the standard model based on the $SU(3)_C \otimes SU(3)_L \otimes U(1)_N$ gauge symmetry can be divided into two classes accordingly if there are two charged [5,6] or two neutral leptons [7] in each leptonic multiplet. In each class there are several versions according to the number of singlets that are added. We consider a typical model of the first class (Model A) and a model of the second class (Model B).

The covariant derivatives of the triplet ϕ are

$$\mathcal{D}_\mu \phi_i = \partial_\mu \phi_i - ig \left(\vec{W}_\mu \cdot \frac{\vec{\lambda}}{2} \right)_i^j \phi_j - ig' N_{\phi_i} B_\mu \phi_i, \quad (1a)$$

and for the symmetric tensor S we have

$$\mathcal{D}_\mu S_{mn} = \partial_\mu S_{mn} - ig \left[\left(\vec{W}_\mu \cdot \frac{\vec{\lambda}}{2} \right)_m^k S_{kn} + \left(\vec{W}_\mu \cdot \frac{\vec{\lambda}}{2} \right)_n^k S_{km} \right] - ig' N B_\mu S_{mn}; \quad (1b)$$

In general, there are several distinct models depending on the way the electric charge operator is embedded in the generators of $SU(3)_L \otimes U(1)_N$. We will use the leptonic sector, with the usual charges, 0 and ± 1 , to define the representation content of several 3-3-1 models.

Model A is defined by the charge operator

$$Q_A = \frac{1}{2} (\lambda_3 - \sqrt{3} \lambda_8) + N, \quad (2a)$$

In this sort of models, besides the quarks of charge $2/3$ and $-1/3$, there are exotic quarks with charges $5/3$ and $-4/3$.

On the other hand, Model B is characterized by the charge operators

$$Q_B = \frac{1}{2} \left(\bar{\lambda}_3 + \frac{1}{\sqrt{3}} \bar{\lambda}_8 \right) + N, \quad (2b)$$

with $\bar{\lambda} = -\lambda^*$. In this kind of models all quarks have charges $2/3$ or $-1/3$.

In both types of models the Spontaneous Symmetry Breakdown (SSB)

$$SU(3)_L \otimes U(1)_N \longrightarrow SU_L(2) \otimes U(1)_Y \longrightarrow U(1)_Q, \quad (3)$$

is obtained by at least two scalar triplets. However, in order to give mass to the quarks a third scalar triplet must be added or, in some cases, for giving a mass to the leptons a scalar sextet is also required. These multiplets contribute to the masses of the charged and neutral vector bosons too.

We have chosen the λ 's matrices in such a way that the first two entries in each triplet or antitriplet are the $SU(2)_L$ particles while the third one may be a new field.

In Model A, Eq. (2a) implies

$$\Psi_{(A)lL} = \begin{pmatrix} \nu_l \\ l^- \\ l^+ \end{pmatrix}_L \sim (\mathbf{3}, 0), \quad l = e, \mu, \tau. \quad (4)$$

Notice that the lepton multiplet in Eq. (4) is that of Ref. [5]. Since we are concerned here with the SSB aspect we will write explicitly only the scalar multiplets. In this model, the introduction of right-handed neutrinos is optional.

Leptons with electric charge 0 and ± 1 in model B are according to Eq. (2b)

$$\Psi_{(B)lL} = \begin{pmatrix} \nu_l \\ l^- \\ \nu_l^c \end{pmatrix}_L \sim (\mathbf{3}, -1/3). \quad (5)$$

In this sort of models, the right-handed charged leptons are in singlets of the gauge symmetry. Neutrinos can, in principle, get a Dirac or Majorana mass, depending on the representation content of the scalar sector and also, on the VEV of the neutral scalar fields implemented by an appropriated scalar potential.

In all these models one of the quark families transforms in the same way as leptons. It is not necessarily the first family which does so. The distinction among families will be necessary only after the diagonalization of the quark mass matrices. Since we are concerned here only with weak eigenstates, we do not need to distinguish which family transforms in

the same way than the leptons. This will be manifested only in the parametrization of the mixing matrices V_L^U , V_L^D and V_R^U , V_R^D . However, we stress that from the phenomenological point of view it will be important to consider if it is the lightest family, or the heaviest one, which has its components formed mainly by those fields transforming in the same way than the leptons.

For model A the scalar triplets are

$$\eta = \begin{pmatrix} \eta^0 \\ \eta_1^- \\ \eta_2^+ \end{pmatrix} \sim (\mathbf{3}, 0), \quad \rho = \begin{pmatrix} \rho^+ \\ \rho^0 \\ \rho^{++} \end{pmatrix} \sim (\mathbf{3}, +1), \quad \chi = \begin{pmatrix} \chi^- \\ \chi^{--} \\ \chi^0 \end{pmatrix} \sim (\mathbf{3}, -1). \quad (6)$$

and the sextet from Eq. (4) is

$$S = \begin{pmatrix} S_{11}^0 & S_{12}^-/\sqrt{2} & S_{13}^+/\sqrt{2} \\ S_{12}^-/\sqrt{2} & S_{22}^{--} & S_{23}^0/\sqrt{2} \\ S_{13}^+/\sqrt{2} & S_{23}^0/\sqrt{2} & S_{33}^{++} \end{pmatrix} \sim (\mathbf{6}, 0). \quad (7)$$

We will use the notation $\langle \eta^0 \rangle \equiv v_\eta$, $\langle \rho^0 \rangle \equiv v_\rho$, $\langle \chi^0 \rangle \equiv v_\chi$, $\langle S_{11}^0 \rangle \equiv v_1$, $\langle S_{23}^0 \rangle \equiv \sqrt{2}v_2$.

From Eqs. (6) and (7) we obtain (up to a $g^2/4$ factor)

$$M_A^2 = \begin{pmatrix} v_\eta^2 + v_\rho^2 + 4v_1^2 + 2v_2^2 & \frac{1}{\sqrt{3}}(v_\eta^2 - v_\rho^2 + 4v_1^2 + 2v_2^2) & -2tv_\rho^2 \\ \frac{1}{\sqrt{3}}(v_\eta^2 - v_\rho^2 + 4v_1^2 + 2v_2^2) & \frac{1}{3}(v_\eta^2 + v_\rho^2 + 4v_\chi^2 + 4v_1^2 + 2v_2^2) & \frac{2t}{\sqrt{3}}(v_\rho^2 + 2v_\chi^2) \\ -2tv_\rho^2 & \frac{2t}{\sqrt{3}}(v_\rho^2 + 2v_\chi^2) & 4t^2(v_\rho^2 + v_\chi^2) \end{pmatrix}. \quad (8)$$

For model B the scalars are η and σ :

$$\eta = \begin{pmatrix} \eta^0 \\ \eta_1^- \\ \eta_2^- \end{pmatrix} \sim (\mathbf{3}, -2/3), \quad \sigma = \begin{pmatrix} \sigma^+ \\ \sigma_1^0 \\ \sigma_2^0 \end{pmatrix} \sim (\mathbf{3}, +1/3). \quad (9)$$

In the present model the sextet has the following charge attribution

$$S = \begin{pmatrix} S_{11}^{++} & S_{12}^+/\sqrt{2} & S_{13}^+/\sqrt{2} \\ S_{12}^+/\sqrt{2} & S_{22}^0 & S_{23}^0/\sqrt{2} \\ S_{13}^+/\sqrt{2} & S_{23}^0/\sqrt{2} & S_{33}^0 \end{pmatrix} \sim (\mathbf{6}, +2/3), \quad (10)$$

with $\langle S_{22}^0 \rangle \equiv v_1$, $\langle S_{33}^0 \rangle \equiv v_2$, $\langle S_{23}^0 \rangle \equiv \sqrt{2}v_3$. We will use the notation $\langle \eta^0 \rangle \equiv v_\eta$, $\langle \sigma_1^0 \rangle \equiv v_{s1}$ and $\langle \sigma_2^0 \rangle \equiv v_{s2}$.

From the Higgs multiplets in Eqs. (9) and (10) we obtain the mass matrix (up to a $g^2/4$ factor)

$$M_B^2 = \begin{pmatrix} m_{11} & \frac{1}{\sqrt{3}}m_{12} & -\frac{8t}{\sqrt{3}}m_{13} \\ \frac{1}{\sqrt{3}}m_{12} & \frac{1}{3}m_{22} & \frac{2t}{3\sqrt{3}}m_{23} \\ -\frac{2t}{\sqrt{3}}m_{13} & \frac{2t}{3\sqrt{3}}m_{23} & \frac{4t^2}{9}m_{33} \end{pmatrix}, \quad (11a)$$

where

$$m_{11} = v_\eta^2 + v_{s1}^2 + 4v_1^2 + 2v_2^2, \quad m_{12} = -v_\eta^2 + v_{s1}^2 + 4v_1^2 - 2v_2^2, \quad (11b)$$

$$m_{13} = v_\eta^2/2 + v_{s1}^2/4 + v_1^2 + 2v_2^2, \quad m_{22} = v_\eta^2 + v_{s1}^2 + 4v_{s2}^2 + 4v_1^2 + 2v_2^2 + 16v_3^2, \quad (11c)$$

$$m_{23} = 2v_\eta^2 - v_{s1}^2 + 2v_{s2}^2 - 4v_1^2 + 4v_2^2 + 8v_3^2, \quad m_{33} = 4v_\eta^2 + v_{s1}^2 + v_{s2}^2 + 4v_1^2 + 8v_2^2 + 4v_3^2. \quad (11d)$$

It is straightforward to verify that both matrices in Eqs. (8) and (11a) are singular *i.e.*, $\det M^2 = 0$ even if all VEVs are different from zero. Hence, contrary to what has been claimed in Refs. [3] and [4] we have the photon massless as usual even if all the neutral components of the sextets in Eqs. (7) and (10) gain a VEV.

In model B without the sextet, neutrino masses come only through Yukawa couplings to the triplet η : $f_{lm}\epsilon^{ijk}\overline{\Psi}_{iL}^c\Psi_{mjL}\eta_k$ being f_{lm} antisymmetric matrix in the flavor space. So, the neutrino mass spectrum is: one is massless and the other two are mass degenerate. However, since neutrinos and antineutrinos are placed in this model in the same multiplet, the mass degeneracy can be broken by radiative corrections, inducing Majorana masses if we choose an appropriate scalar potential [8]. Or, we can make $\nu_L^c \rightarrow \mathcal{N}_L^c$ in Eq. (5), being $\mathcal{N}$'s heavy neutral leptons and also adding the respective right-handed components ν_R 's and $\mathcal{N}_R^c$'s. In this situation all neutral leptons gain an arbitrary Dirac mass via the Yukawa interaction with the triplets. This also happens in model A which include heavy charged leptons [9].

Summarizing, in both kind of models it is possible all the VEVs of the sextet be non-zero giving mass to the charged or neutral leptons and there is not a relation between any of the VEVs of those sextets and the breaking of the electromagnetic gauge invariance as claimed in both Refs. [3] and [4] and some time before in Ref. [7]. We have verified also that this does not depend either on the specific model. For instance the model of Ref. [10] is equivalent to model A and the same results are obtained: the charge operators can be related by a unitary transformation. In fact, by using the $SU(3)$ matrix

$$U = \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (12)$$

the operator Q_A in Eq. (2a) turns into the charge operator defined in Ref. [10]. Similar equivalence occurs in models of the type of Model B that have been treated in literature as in Ref. [11].

ACKNOWLEDGMENTS

We thank (V.P.) Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for partial financial support and IF-UERJ and IFT-UNESP (M.D.T.) for hospitality.

REFERENCES

- [1] L-F. Li, Phys. Rev. **D9** (1974) 1723.
- [2] J. D. Bjorken and C. H. Llewellyn Smith, Phys. Rev. D **7** (1973) 887.
- [3] M. Özer, Phys. Lett. **B 337** (1994) 324.
- [4] F. Pisano, J.A. Silva-Sobrinho and M. D. Tonasse, Phys. Lett. **B 388** (1996) 338.
- [5] F. Pisano and V. Pleitez, Phys. Rev. D **46** (1992) 410.
- [6] R. Foot, O. F. Hernandez, F. Pisano and V. Pleitez, Phys. Rev. D **47** (1993) 4158.
- [7] J. C. Montero, F. Pisano and V. Pleitez, Phys. Rev. D **47** (1993) 2918.
- [8] R. Barbieri and R. Mohapatra, Phys. Lett. **B218** (1989) 225.
- [9] V. Pleitez and M. D. Tonasse, Phys. Rev. D **48** (1993) 2353.
- [10] P. H. Frampton, Phys. Rev. Lett. **69** (1992) 2889.
- [11] J. Liu, Phys. Lett. **B 225** (1989) 148 and references therein.